arXiv:1410.1483v1 [math.GR] 6 Oct 2014

On t-extensions of abelian groups

H.Sahleh
Department of Mathematics, University of Guilan, P.O.BOX 1914
Rasht-Iran
e-mail: sahleh@guilan.ac.ir
A.A. Alijani
Department of Mathematics, University of Guilan
e-mail: taleshalijan@phd.guilan.ac.ir

Abstract

Let $\Re$ be the category of all discrete abelian groups and $\pounds$ the category of all locally compact abelian (LCA) groups. For a group $G \in \pounds$, the maximal torsion subgroup of G is denoted by tG. A short exact sequence $0 \to A \xrightarrow{\phi} B \xrightarrow{\psi} C \to 0$ in $\Re$ is said to be a t-extension if $0 \to tA \xrightarrow{\phi} tB \xrightarrow{\psi} tC \to 0$ is a short exact sequence. We show that the set of all t-extensions of A by C is a subgroup of $Ext(C, A)$ which contains $Pext(C, A)$ for discrete abelian groups A and C. We establish conditions under which the t-extensions split and determine those groups in $\Re$ which are t-injective or t-projective in $\Re$. Finally we determine the compact groups G in $\pounds$ such that every pure extension $0 \to G \to Y \to C \to 0$ splits where C is a compact connected group in $\pounds$.

Introduction

Throughout, all groups are Hausdorff topological abelian groups and will be written additively. Let $\pounds$ denote the category of locally compact abelian (LCA) groups with continuous homomorphisms as morphisms and $\Re$ the category of discrete abelian groups. The Pontrjagin dual and the maximal torsion subgroup of a group $G \in \pounds$ are denoted by $\hat{G}$ and tG,respectively. A morphism is called proper if it is open onto its image and a short exact

sequence $0 \to A \xrightarrow{\phi} B \xrightarrow{\psi} C \to 0$ in $\pounds$ is said to be proper exact if ϕ and ψ are proper morphisms. In this case the sequence is called an extension of A by C (in $\pounds$). Following [4], we let $Ext(C, A)$ denote the (discrete) group of extensions of A by C. A subgroup H of a group C is called pure if $nH = H \cap nC$ for all positive integers n. An extension $0 \to A \xrightarrow{\phi} B \xrightarrow{\psi} C \to 0$ is called a pure extension if $\phi(A)$ is pure in B. The elements represented by pure extensions of A by C form a subgroup of $Ext(C, A)$ which is denoted by $Pext(C, A)$. In section 1 and 2, all groups are discrete abelian groups. An extension $0 \to A \xrightarrow{\phi} B \xrightarrow{\psi} C \to 0$ in $\Re$ will be called a t-extension if $0 \to tA \xrightarrow{\phi} tB \xrightarrow{\psi} tC \to 0$ is an extension. Let $Ext_t(C, A)$ denote the set of all elements in $Ext(C, A)$ represented by t-extensions. In section 1, we show that $Ext_t(C, A)$ is a subgroup of $Ext(C, A)$ which contains $Pext(C, A)$ (see Theorem 1.6 and Lemma 1.7). In section 2, we establish some results on splitting of t-extensions (see Lemma 2.1, Theorem 2.11 and Theorem 2.13). Assume that $\Im$ is any subcategory of $\pounds$. The section four is a part of an investigation which answers the following question:
Under what conditions on $G \in \pounds$, $Ext(X, G) = 0$ or $Pext(X, G) = 0$ for all $X \in \Im$?. In [2], [3], [4], [5], [7] and [8] the question is answered in some subcategories of $\pounds$ such as the category of divisible locally compact abelian groups. In [5, Corollary 3.4], Fulp and Griffith proved that a compact group G satisfies $Ext(C, G) = 0$ for all compact connected groups C if and only if $G \cong (R/Z)^\sigma$ where σ is a cardinal. It may happen that $Ext(X, G) \neq 0$ but $Pext(X, G) = 0$. For example, $Ext(Z_n, Z) \neq 0$ but $Pext(Z_n, Z) = 0$ where Z is the group of integers and Z_n is the cyclic group of order n. In this paper, we show that a compact group G satisfies $Pext(C, G) = 0$ for all compact connected groups C if and only if $G \cong (R/Z)^\sigma \bigoplus H$ where H is a compact totally disconnected group (see Theorem 3.2). For the characterization of compact groups G which $Pext(C, G) = 0$ for all compact connected groups C, we need to show that $Pext(X, A) = 0$ for a discrete torsion group X and a discrete torsion-free group A (see Corollary 2.2). As a result,[5, Corollary 3.4] is a consequence of Theorem 3.2 (see Remark 3.3).

The additive topological group of real numbers is denoted by R and Q is the group of rationals with the discrete topology. We denote by G_0, the identity component of a group $G \in \pounds$. For more on locally compact abelian groups, see [6].

1 t-extension

In this section, we define the concept of a t-extension of A by C. We show that the set of all t-extensions of A by C form a subgroup of $Ext(C, A)$ which contains $Pext(C, A)$.

Definition 1.1. *An extension* $0 \to A \xrightarrow{\phi} B \xrightarrow{\psi} C \to 0$ *is called a t-extension if* $0 \to tA \xrightarrow{\phi} tB \xrightarrow{\psi} tC \to 0$ *is an extension.*

Lemma 1.2. *A pushout or a pullback of a t-extension is a t-extension.*

Proof. Suppose $0 \to A \xrightarrow{\phi} B \xrightarrow{\psi} C \to 0$ is a t-extension and

$$\begin{array}{ccccccccc} 0 & \longrightarrow & A & \xrightarrow{\phi} & B & \xrightarrow{\psi} & C & \longrightarrow & 0 \\ & & \downarrow{\scriptstyle \mu} & & \downarrow & & \downarrow{\scriptstyle 1_C} & & \\ 0 & \longrightarrow & A' & \xrightarrow{\phi'} & (A' \bigoplus B)/H & \xrightarrow{\psi'} & C & \longrightarrow & 0 \end{array}$$

is a standard pushout diagram (see [1]). Then

$$H = \{(\mu(a), -\phi(a)), a \in A\}$$

and

$$\phi' : a' \longmapsto (a', 0) + H, \qquad \psi' : (a', b) + H \longmapsto \psi(b)$$

We show that $0 \to tA' \xrightarrow{\phi'} t((A' \bigoplus B)/H) \xrightarrow{\psi'} tC \to 0$ is exact. First, we show that $\psi' : t((A' \bigoplus B)/H) \to tC$ is surjective. Let $c \in tC$. Since $0 \to tA \xrightarrow{\phi} tB \xrightarrow{\psi} tC \to 0$ is exact, so there exists $b \in tB$ such that $\psi(b) = c$. Clearly, $(0, b) + H \in t((A' \bigoplus B)/H)$. On the other hand, $\psi'((0, b) + H) = \psi(b) = c$. Hence ψ' is surjective. Now, we show that $Ker\psi' \mid_X \subseteq Im\phi' \mid_{tA'}$ where $X = t((A' \bigoplus B)/H)$. Let $(a', b) + H \in X$ and $\psi'((a', b) + H) = 0$. So, $\psi(b) = 0$. Hence, there exists $a \in A$ such that $\phi(a) = -b$. On the other hand, there exists a positive integer n such that $(na', nb) \in H$. So, there exists $a_1 \in A$ such that $\mu(a_1) = na'$ and $-\phi(a_1) = nb$. Now, we have

$$\phi(a_1 - na) = \phi(a_1) - n\phi(a) = 0$$

So $a_1 = na$ and $n(a' - \mu(a)) = 0$. It follows that $a' - \mu(a) \in tA'$ and $\phi'(a' - \mu(a)) = (a' - \mu(a), 0) + H = (a', b) + H$ (since $(a' - \mu(a), 0) - (a', b) =$

$(-\mu(a), -b) = (\mu(-a), -\phi(-a)) \in H)$. Now, suppose that

$$\begin{array}{ccccccccc} 0 & \longrightarrow & A & \xrightarrow{\phi'} & B' & \xrightarrow{\psi'} & C' & \longrightarrow & 0 \\ & & \downarrow{\scriptstyle 1_A} & & \downarrow & & \downarrow{\scriptstyle \gamma} & & \\ 0 & \longrightarrow & A & \xrightarrow{\phi} & B & \xrightarrow{\psi} & C & \longrightarrow & 0 \end{array}$$

is a standard pullback diagram. Then

$$B' = \{(b, c'); \psi(b) = \gamma(c')\}$$

and

$$\phi' : a \longmapsto (\phi(a), 0), \qquad \psi' : (b, c') \longmapsto c'$$

We show that $0 \to tA \xrightarrow{\phi'} tB' \xrightarrow{\psi'} tC' \to 0$ is exact. Let $c' \in tC'$. Then, there exists a positive integer n such that $nc' = 0$. Since ψ is surjective, $\psi(b) = \gamma(c')$ for some $b \in B$. Now, $n\psi(b) = \gamma(nc') = 0$. Hence, $\psi(b) \in tC$. Since $0 \to tA \xrightarrow{\phi} tB \xrightarrow{\psi} tC \to 0$ is exact, so $\psi(b_1) = \psi(b)$ for some $b_1 \in tB$. Hence, $(b_1, c') \in tB'$ and $\psi'(b_1, c') = c'$. Therefore, $\psi' : tB' \to tC'$ is surjective. Now, suppose that $(b, c') \in tB'$ and $\psi'(b, c') = 0$. Then $c' = 0$ and $nb = 0$ for some positive integer n. So $b \in tB$. Since $\psi(b) = \gamma(c') = 0$ and $0 \to tA \xrightarrow{\phi} tB \xrightarrow{\psi} tC \to 0$ is exact, there exists $a \in tA$ such that $\phi(a) = b$. Now, we have

$$\phi'(a) = (\phi(a), 0) = (b, 0) = (b, c')$$

It follows that $Ker\psi' \mid_{tB'} \subseteq Im\phi' \mid_{tA}$.

Remark 1.3. Let $\beta : B \to X$ be an isomorphism and $x \in tX$. Then $nx = 0$ for some positive integer n. Since β is surjective, so there exist $b \in B$ such that $\beta(b) = x$. Hence, $\beta(nb) = 0$. Since β is injective, so $nb = 0$. Therefore, $\beta \mid_{tB}: tB \to tX$ is an isomorphism.

Recall that two extensions $0 \to A \xrightarrow{\phi_1} B \xrightarrow{\psi_1} C \to 0$ and $0 \to A \xrightarrow{\phi_2} X \xrightarrow{\psi_2} C \to 0$ are said to be equivalent if there is an isomorphism $\beta : B \to X$ such that the following diagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & A & \xrightarrow{\phi_1} & B & \xrightarrow{\psi_1} & C & \longrightarrow & 0 \\ & & \downarrow{\scriptstyle 1_A} & & \downarrow{\scriptstyle \beta} & & \downarrow{\scriptstyle 1_C} & & \\ 0 & \longrightarrow & A & \xrightarrow{\phi_2} & X & \xrightarrow{\psi_2} & C & \longrightarrow & 0 \end{array}$$

is commutative.

Lemma 1.4. An extension equivalent to a t-extension is a t-extension.

Proof. Let

$$E_1 : 0 \to A \xrightarrow{\phi_1} B \xrightarrow{\psi_1} C \to 0$$

and

$$E_2 : 0 \to A \xrightarrow{\phi_2} X \xrightarrow{\psi_2} C \to 0$$

be two equivalent extensions such that E_1 is a t-extension. Then, there is an isomorphism $\beta : B \to X$ such that $\beta\phi_1 = \phi_2$ and $\psi_2\beta = \psi_1$. Let $x \in tC$. Since E_1 is t-extension, so $\psi_1(b) = x$ for some $b \in tB$. Hence, $\psi_2(\beta(b)) = \psi_1(b) = x$. So, $\psi_2 : tX \to tC$ is surjective. Now, let $\psi_2(x) = 0$ for some $x \in tX$. By Remark 1.3, there exist $b \in tB$ such that $\beta(b) = x$. Hence, $\psi_1(b) = \psi_2(\beta(b)) = 0$. Since E_1 is t-extension, so $\phi_1(a) = b$ for some $a \in tA$. Consequently, $\phi_2(a) = \beta(\phi_1(a)) = x$.

Remark 1.5. Let C and A be two groups and $0 \to A \xrightarrow{\phi_1} B_1 \xrightarrow{\psi_1} C \to 0$ and $0 \to A \xrightarrow{\phi_2} B_2 \xrightarrow{\psi_2} C \to 0$ two t-extensions of A by C. An easy calculation shows that $0 \to A \bigoplus A \xrightarrow{(\phi_1 \bigoplus \phi_2)} B_1 \bigoplus B_2 \xrightarrow{(\psi_1 \bigoplus \psi_2)} C \bigoplus C \to 0$ is a t-extension where $(\phi_1 \bigoplus \phi_2)(a_1, a_2) = (\phi_1(a_1), \phi_2(a_2))$ and $(\psi_1 \bigoplus \psi_2)(b_1, b_2) = (\psi_1(b_1), \psi_2(b_2))$.

Theorem 1.6. Let A and C be two groups. Then, the class $Ext_t(C, A)$ of all equivalence classes of t-extensions of A by C is an subgroup of $Ext(C, A)$ with respect to the operation defined by

$$[E_1] + [E_2] = [\nabla_A(E_1 \bigoplus E_2)\triangle_C]$$

where E_1 and E_2 are t-extensions of A by C and ∇_A and $\triangle_C$ are the diagonal and codiagonal homomorphism.

Proof. Clearly, $0 \to A \to A \bigoplus C \to C \to 0$ is a t-extension. By Lemma 1.2 and Remark 1.5, $[E_1] + [E_2] \in Ext_t(C, A)$ for two t-extensions E_1 and E_2 of A by C. So, $Ext_t(C, A)$ is a subgroup of $Ext(C, A)$.

Lemma 1.7. *Let A and C be two groups. Then, $Pext(C, A) \subseteq Ext_t(C, A)$.*

Proof. Let $0 \to A \xrightarrow{\phi} B \xrightarrow{\psi} C \to 0$ be an element of $Pext(C, A)$. It is sufficient to show that $tB/t\phi(A) \cong t(B/\phi(A))$. Consider the map $\varphi : tB \to t(B/\phi(A))$ given by $b \longmapsto b+\phi(A)$. Clearly, φ is a homomorphism. We show that φ is surjective. Let $b+\phi(A) \in t(B/\phi(A))$. Then, there exists a positive integer n such that $nb \in \phi(A)$. Since $\phi(A)$ is pure in B, so $nb = n\phi(a)$ for some $a \in A$. Hence, $n(b-\phi(a)) = 0$. This shows that $b-\phi(a) \in tB$ and $\varphi(b-\phi(a)) = b+\phi(A)$. So φ is surjective. We have

$$Ker\varphi = \{b \in tB; b \in \phi(A)\} = \phi(A) \bigcap tB = t\phi(A)$$

Hence $tB/t\phi(A) \cong t(B/\phi(A))$.

Corollary 1.8. *If A is a divisible group or C a torsion-free group, then $Pext(C, A) = Ext_t(C, A) = Ext(C, A)$.*

Proof. It is clear.

2 Splitting of t-extensions

In this section, we establish some conditions on A and C such that $Ext_t(C, A) = 0$. We also determine the t-injective and t-projective groups in $\Re$.

Lemma 2.1. *Let A be a torsion-free group and C a torsion group. Then, $Ext_t(C, A) = 0$.*

Proof. Let $E : 0 \to A \xrightarrow{\phi} B \xrightarrow{\psi} C \to 0$ be a t-extension. Then $\psi_{|tB} : tB \to C$ is an isomorphism. Let $b \in B$. Then, $\psi(b) \in C$. So $\psi(b) = \psi(b')$ for some $b' \in tB$. Hence, $b - b' = \phi(a)$ for some $a \in A$. This follows that $B = \phi(A) + tB$. Since $\phi(A)$ is torsion-free and tB torsion, so $\phi(A) \bigcap tB = 0$. Hence, $B = \phi(A) \bigoplus tB$ and E splits.

Corollary 2.2. *Let A be a torsion-free group and C a torsion group. Then, $Pext(C, A) = 0$.*

Proof. It is clear by Lemma 1.7 and Lemma 2.1.

Lemma 2.3. Let A and C be two torsion groups. Then $Ext(C,A) = Ext_t(C,A)$.

Proof. Let A and C be two torsion groups. It is clear that $Ext_t(C,A) \subseteq Ext(C,A)$. Suppose that $E : 0 \to A \to B \to C \to 0$ is an extension. Then, B is a torsion group. Hence, E is a t-extension.

Lemma 2.4. *Let C be a torsion group. Then, $Ext(C,tA) \cong Ext_t(C,A)$ for every group A.*

Proof. The exact sequence $0 \to tA \xrightarrow{i} A \xrightarrow{\pi} A/tA \to 0$ induces the following exact sequence

$$Hom(C,A/tA) \to Ext(C,tA) \xrightarrow{i_*} Ext(C,A) \xrightarrow{\pi_*} Ext(C,A/tA) \to 0$$

Note that $Hom(C,A/tA) = 0$. By Lemma 2.1, $Ext_t(C,A/tA) = 0$. Since $\pi_*(Ext_t(C,A)) \subseteq Ext_t(C,A/tA)$, so $\pi_*(Ext_t(C,A)) = 0$. Hence, $Ext_t(C,A) \subseteq Ker\pi_* = i_*(Ext(C,tA))$. By Lemma 2.3, $Ext(C,tA) = Ext_t(C,tA)$. So, $i_*(Ext(C,tA)) \subseteq Ext_t(C,A)$. Hence, $Ext(C,tA) \cong Ext_t(C,A)$.

Corollary 2.5. *Let A be a group. Then, $Ext_t(Z(m),A) \cong tA/m(tA)$ for every positive integer m.*

Proof. It is clear by Lemma 2.4 and [1,page 222].

Corollary 2.6. *Let C be a torsion group and $\{A_i : i \in I\}$ a collection of groups. If I is finite, then $Ext_t(C, \Pi_{i\in I} A_i) \cong \Pi_{i\in I} Ext_t(C,A_i)$.*

Proof. Let $I = \{1,...,n\}$ for some positive integer n. By Lemma 2.4, $Ext_t(C, \Pi_{i=1}^n A_i) \cong Ext(C, t(\Pi_{i=1}^n A_i))$. Since $t(\Pi_{i=1}^n A_i) = \Pi_{i=1}^n t(A_i)$, so $Ext_t(C, \Pi_{i=1}^n A_i) \cong \Pi_{i\in I} Ext(C,tA_i) \cong \Pi_{i\in I} Ext_t(C,A_i)$.

Remark 2.7. *In general, $Ext_t(C, \Pi_{i\in I} A_i) \ncong \Pi_{i\in I} Ext_t(C,A_i)$.*

Example 2.8. *Let p be a prime and $H = \prod_{n=1}^{\infty} Z(p^n)$. By Lemma 2.4, $Ext_t(Q/Z,H) \cong Ext(Q/Z,tH)$. Consider the following exact sequence*

$$0 \to Ext(Q/Z,tH) \to Ext(Q/Z,H) \to Ext(Q/Z,H/tH) \to 0 \quad (2.1)$$

By [1, Theorem 52.2] and Lemma 2.3,

$$Ext(Q/Z, H) \cong \prod_{n=1}^{\infty} Ext(Q/Z, Z(p^n)) \cong \prod_{n=1}^{\infty} Ext_t(Q/Z, Z(p^n))$$

If $Ext_t(Q/Z, H) \cong \prod_{n=1}^{\infty} Ext_t(Q/Z, Z(p^n))$,*then* $Ext(Q/Z, H) \cong Ext(Q/Z, tH)$. *It follows from (2.1) that,* $Ext(Q/Z, H/tH) = 0$ *which is a contradiction, since* H/tH *is not divisible.*

Lemma 2.9. *Let* A *be a torsion-free group. Then,* $Ext(C/tC, A) \cong Ext_t(C, A)$ *for every group* C.

Proof. The exact sequence $0 \to tC \xrightarrow{i} C \xrightarrow{\pi} C/tC \to 0$ *induces the following exact sequence*

$$Hom(tC, A) \to Ext(C/tC, A) \xrightarrow{\pi_*} Ext(C, A) \xrightarrow{i_*} Ext(tC, A) \to 0$$

Note that $Hom(tC, A) = 0$. *By Lemma 2.1,* $Ext_t(tC, A) = 0$. *So,* $i_*(Ext_t(C, A)) \subseteq Ext_t(tC, A) = 0$. *Hence,* $Ext_t(C, A) \subseteq Keri_* = \pi_*(Ext(C/tC, A))$. *By Corollary 1.8,* $Ext(C/tC, A) = Ext_t(C/tC, A)$. *So,* $\pi_*(Ext(C/tC, A)) \subseteq Ext_t(C, A)$. *Hence,* $Ext(C/tC, tA) \cong Ext_t(C, A)$.

Definition 2.10. *Let* G *be a group. We call* G *a t-injective group in* $\Re$ *if for every t-extension*

$$0 \to A \xrightarrow{\phi} B \to C \to 0$$

and a homomorphism $f : A \to G$, *there is a homomorphism* $\bar{f} : B \longrightarrow G$ *such that* $\bar{f}\phi = f$.

We call G *a t-projective group in* $\Re$ *if for every t-extension*

$$0 \to A \to B \xrightarrow{\psi} C \to 0$$

and a homomorphism $f : G \to C$, *there is a homomorphism* $\bar{f} : G \to B$ *such that* $\psi\bar{f} = f$.

Recall that a group A *is said to be cotorsion if* $Ext(Q, A) = 0$.

Theorem 2.11. *Let* A *be a group. The following statements are equivalent:*

1. *A is t-injective in $\Re$.*

2. *$Ext_t(C, A) = 0$ for all $C \in \Re$.*

3. *$A \cong B \bigoplus D$ where B is a torsion divisible group and D a torsion-free cotorsion group.*

Proof. $1 \Longrightarrow 2$*: Let A be a t-injective in $\Re$ and $E : 0 \to A \xrightarrow{\phi} B \longrightarrow C \to 0$ a t-extension of A by C. Then, there is a homomorphism $\bar{\phi} : B \to G$ such that $\bar{\phi}\phi = 1_G$. Consequently, E splits.*

$2 \Longrightarrow 3$*: Let $Ext_t(C, A) = 0$ for every group C. So $Ext_t(Z(m), A) = 0$ for every positive integer m. By Corollary 2.5, $m(tA) = tA$ for every positive integer m. So, tA is divisible. Hence, $A \cong tA \bigoplus A/tA$. Therefore, $Ext(Q, A) \cong Ext(Q, tA) \bigoplus Ext(Q, A/tA)$. Since $Ext(Q, A) = Ext_t(Q, A) = 0$, then $Ext(Q, A/tA) = 0$. Hence, A/tA is cotorsion. Now, we set $tA = B$ and $A/tA = D$.*

$3 \Longrightarrow 2$*: Suppose that $A \cong B \bigoplus D$ where B is a torsion divisible group and D a torsion-free cotorsion group. Let C be a group. Since $Ext(C, B) = 0$, so $\pi_2^* : Ext(C, A) \to Ext(C, D)$ is an isomorphism. By Lemma 2.9, $Ext_t(C, D) \cong Ext(C/tC, D)$. Since D is a cotorsion group, so $Ext(C/tC, D) = 0$. Hence, $Ext_t(C, D) = 0$. So, $\pi_2^*(Ext_t(C, A)) \subseteq Ext_t(C, D) = 0$. Since π_2^* is an isomorphism, so $Ext_t(C, A) = 0$.*

$2 \Longrightarrow 1$*: Let $E : 0 \to G \xrightarrow{\phi} B \to C \to 0$ be a t-extension and $f : G \to A$ a homomorphism. Then f induces a pushout diagram*

$$\begin{array}{ccccccccc} E: 0 & \longrightarrow & G & \xrightarrow{\phi} & B & \longrightarrow & C & \longrightarrow & 0 \\ & & \downarrow{\scriptstyle f} & & \downarrow & & \downarrow & & \\ fE: 0 & \longrightarrow & A & \xrightarrow{\mu} & (A \bigoplus B)/H & \longrightarrow & C & \longrightarrow & 0 \end{array}$$

Where $H = \{(-f(a), \phi(a)); a \in A\}$ and $\mu : a \longmapsto (a, 0) + H$. By Lemma 1.2, fE is a t-extension and by assumption it splits. Hence A is t-injective.

Recall that a group A is called algebraically compact if and only if $Pext(X, A) = 0$ for every group X [1].

Corollary 2.12. *A torsion-free, cotorsion group is algebraically compact.*

Proof. *Let A be a torsion-free, cotorsion group. By Theorem 2.11, $Ext_t(C, A) = 0$ for every group C. Hence, $Pext(C, A) = 0$ for every group C.*

Theorem 2.13. *Let C be a group. The following statements are equivalent:*

1. *C is t-projective in $\Re$.*
2. *$Ext_t(C, A) = 0$ for all $A \in \Re$.*
3. *C is a free group.*

Proof. *$1 \Longrightarrow 2$: Let C be a t-projective in $\Re$ and $E : 0 \to A \xrightarrow{\phi} B \longrightarrow C \to 0$ a t-extension of A by C. Then there is a homomorphism $\bar{\phi} : C \to B$ such that $\bar{\phi}\phi = 1_C$. Consequently, E splits.*

$2 \Longrightarrow 3$: Let $Ext_t(C, A) = 0$ for every group A. By Lemma 1.7, $Pext(C, A) = 0$ for every group A. Hence, C is free group.

$3 \Longrightarrow 1$: It is clear.

3 Splitting of pure extensions in the category of locally compact abelian groups

In this section, we determine the structure of a compact group G such that $Pext(C, G) = 0$ for all compact connected groups C.

Lemma 3.1. *If A is a compact totally disconnected group and C a compact connected group, then $Pext(C, A) = 0$.*

Proof. *By [7, Lemma 2.3], $Pext(C, A) \cong Pext(\hat{A}, \hat{C})$. On the other hand, $\hat{A}$ and $\hat{C}$ are a discrete, torsion group and a discrete, torsion-free respectively [6, Theorem 24.25 and 24.26]. Hence, by Corollary 2.2,$Pext(\hat{A}, \hat{C}) = 0$. So $Pext(C, A) = 0$.*

Let G be a compact group. Then, $Ext(C, G) = 0$ for every compact connected group C if and only if $G \cong (R/Z)^\sigma$ [5, Corollary 3.4]. In the next Theorem, we show that $Pext(C, G) = 0$ for every compact connected group

C if and only if $G \cong (R/Z)^\sigma \bigoplus H$ where H is a compact totally disconnected group.

Theorem 3.2. *Let G be a compact group. Then, $Pext(C, G) = 0$ for all compact connected groups C if and only if $G \cong (R/Z)^\sigma \bigoplus H$ where H is a compact totally disconnected group.*

Proof. Let G be a compact group and C a compact connected group. Consider the exact sequence $0 \to G_0 \to G \to G/G_0$. By Proposition 4 of [2], we have the exact sequence

$$Hom(C, G/G_0) \to Ext(C, G_0) \to Pext(C, G) \to Pext(C, G/G_0)$$

By Lemma 3.1, $Pext(C, G/G_0) = 0$. Since G/G_0 is totally disconnected, so $Hom(C, G/G_0) = 0$. It follows that $Pext(C, G) \cong Ext(C, G_0)$. Now, Let $Pext(C, G) = 0$ for all compact connected groups C. Then $Ext(C, G_0) = 0$ for all compact connected groups C. By [5, Corollary 3.4], $G_0 \cong (R/Z)^\sigma$. So, the extension $0 \to G_0 \to G \to G/G_0 \to 0$ splits. This shows that $G \cong (R/Z)^\sigma \bigoplus G/G_0$. The converse is clear.

Remark 3.3. *Let G be a compact group such that $Ext(C, G) = 0$ for every compact connected group C. Then $Pext(C, G) = 0$ for every compact connected group C. So by Theorem 3.2, $G \cong (R/Z)^\sigma \bigoplus H$ where H is a compact totally disconnected group. Since R/Z is a compact connected group, so $Ext(R/Z, H) = 0$. Consider the following exact sequence*

$$Hom(R, H) \to Hom(Z, H) \to Ext(R/Z, H) = 0$$

Since R is a connected group and H a totally disconnected group, so $Hom(R, H) = 0$. Hence $H = 0$ and $G \cong R/Z$.

References

[1] *L.Fuchs, Infinite Abelian Groups, Vol. I, Academic Press, New York, 1970.*

[2] *R. O. Fulp, Homological study of purity in locally compact groups, Proc. London Math. Soc. 21 (1970), 501-512.*

[3] R.O.Fulp, Splitting locally compact abelian groups, Michigan Math. J. 19 (1972), 47-55.

[4] R.O.Fulp and P. Griffith, Extensions of locally compact abelian groups I,Trans. Amer. Math. Soc. 154 (1971), 341-356.

[5] R.O.Fulp and P.Griffith, Extensions of locally compact abelian groups II, Trans Math. Soc. 154 (1971), 357-363.

[6] E. Hewitt and K. Ross, Abstract Harmonic Analysis, Vol I, Second Edition, Springer Verlag, Berlin, 1979.

[7] P. Loth, Pure extensions of locally compact abelian groups, Rend.Sem.Mat.Univ.Padova . 116 (2006), 31-40.

[8] H. Sahleh and A. A. Alijani, Splitting of extensions in the category of locally compact abelian groups, International Journal of Group Theory, Vol. 3, No. 3(2014), 39-45.